

Reproduced by

**Armed Services Technical Information Agen**  
**DOCUMENT SERVICE CENTER**

**KNOTT BUILDING, DAYTON, 2, OHIO**

**AD -**

**17504**

**UNCLASSIFIED**

**Technical Report No. 4**

**THE FINITE STURM-LIOUVILLE TRANSFORM**

**By**

**A. Cemal Eringen**

**to**

**Office of Naval Research  
Department of the Navy**

**Contract N7onr-32909**

**Department of Mechanics  
Illinois Institute of Technology  
Technology Center  
Chicago, Illinois**

**1 August 1953**

**AD No. 17504  
ASTIA FILE COPY**

Acknowledgment

The author is deeply indebted  
to Mr. Thomas King for checking the  
analysis.

## TABLE OF CONTENTS

	Page
Acknowledgment . . . . .	1
Table of Contents . . . . .	11
1. Introduction. . . . .	1
2. The Sturm-Liouville Expansion . . . . .	2
3. Finite Sturm-Liouville Transform . . . . .	4
4. Transform of $Lf$ . . . . .	5
5. Solution of Some Partial Differential Equations . . . . .	6
6. Solutions $\phi_0$ and $\chi_0$ of Some Special Equations . . . . .	9
7. Application . . . . .	11
References . . . . .	17

# The Finite Sturm-Liouville Transform

By

A. Cemal Eringen  
Illinois Institute of Technology

## 1. Introduction

It is known that certain partial differential equations can be solved with the use of particular types of definite integrals having appropriate kernels. The choice of kernel depends on the type of boundary value problem. The solution obtained by these transforms is direct in the sense that it contains the boundary values in the solution. This, of course, is lacking in the classical approach. Finite Fourier transforms are of this type [1].

Recently Tranter [2], used a Legendre polynomial as a kernel. Scott [3] following a similar approach used a Jacobi polynomial as a kernel which extends the result of [2].

It is the purpose of the present paper to extend and unify all such special transforms. Thus we employ a kernel which may be determined to suit each particular type of problem. The transform can be employed to solve a wide class of linear second order partial differential equations.

In this paper we deal only with those transforms whose intervals are finite. Thus, the results obtained here are particularly useful for finite domains, or domains which are finite in one direction. An extension to infinite domains and singular cases will be made in a later paper.

## 2. The Sturm-Liouville Expansion

A second order, linear, homogeneous differential equation containing an arbitrary parameter  $\lambda$  has the general form:

$$Mv = p_0 v_{,yy} + p_1 v_{,y} + (p_2 + \lambda p_3)v = 0 \quad (1)$$

$$a_0 \leq y \leq b_0$$

where  $p_1 = p_1(y)$ ;  $v$  and  $y$  are the dependent and real independent variables, respectively. Indices after comma represent differentiation, i.e.:  $\phi_{,y} = d\phi/dy$ .

Equation (1) can be transformed into the canonical form:

$$Lu = \lambda u, \quad L = q(x) - \frac{d^2}{dx^2}, \quad a \leq x \leq b \quad (2)$$

where:

$$\left. \begin{aligned} u &= \theta(x)v, \quad x = \int (p_3/p_0)^{1/2} dy \\ a &= \int^{a_0} (p_3/p_0)^{1/2} dy, \quad b = \int^{b_0} (p_3/p_0)^{1/2} dy \\ q(x) &= \frac{\theta_{,xx}(x)}{\theta(x)} - \phi(x), \quad \theta(x) = (p_3 k^2/p_0)^{1/4} \\ \phi(x) &= p_2/p_3, \quad k = \exp \int (p_1/p_0) dy \end{aligned} \right\} \quad (3)$$

It is simpler to work with (2).

We consider two solutions  $\phi(x, \lambda)$  and  $\chi(x, \lambda)$  of (2) satisfying the boundary conditions:

$$\begin{aligned} \phi(a, \lambda) &= \sin \alpha, \quad \phi_{,x}(a, \lambda) = -\cos \alpha \\ \chi(b, \lambda) &= \sin \beta, \quad \chi_{,x}(b, \lambda) = -\cos \beta \end{aligned} \quad (4)$$

Both solutions are unique if  $q(x)$  is real, continuous everywhere in  $(a, b)$  and has finite limits for  $x = a$  and  $x = b$

(see [4], p.6). We can build up  $\phi(x, \lambda)$  and  $\chi(x, \lambda)$  as a linear combination of two independent solutions  $\phi_0(x, \lambda)$  and  $\chi_0(x, \lambda)$  of (2). For let:

$$\left. \begin{aligned} \omega_0(\lambda)\phi(x, \lambda) &= \phi_0(x, \lambda)[\chi_0(a, \lambda)\cos\alpha + \chi_{0,x}(a, \lambda)\sin\alpha] \\ &\quad - \chi_0(x, \lambda)[\phi_0(a, \lambda)\cos\alpha + \phi_{0,x}(a, \lambda)\sin\alpha] \\ \omega_0(\lambda)\chi(x, \lambda) &= \phi_0(x, \lambda)[\chi_0(b, \lambda)\cos\beta + \chi_{0,x}(b, \lambda)\sin\beta] \\ &\quad - \chi_0(x, \lambda)[\phi_0(b, \lambda)\cos\beta + \phi_{0,x}(b, \lambda)\sin\beta] \\ \omega_0(\lambda) &= W(\phi_0, \chi_0) = \phi_0(x, \lambda)\chi_{0,x}(x, \lambda) - \phi_{0,x}(x, \lambda)\chi_0(x, \lambda) \end{aligned} \right\} (5)$$

Functions  $\phi(x, \lambda)$  and  $\chi(x, \lambda)$  defined by (5) satisfy (2) and (4). Here  $W(u, v)$  is the Wronskian, and can be shown to be independent of  $x$ .

Let  $\lambda_n$  be a root of  $W(\phi, \chi) = 0$ . Then we have:

$$\omega(\lambda_n) = W(\phi_n, \chi_n) = \phi_n \chi_{n,x} - \phi_{n,x} \chi_n = 0$$

or

$$\phi_{n,x}/\phi_n = \chi_{n,x}/\chi_n$$

where:

$$\phi_n = \phi(x, \lambda_n), \quad \text{and} \quad \chi_n = \chi(x, \lambda_n).$$

Hence the integration gives:

$$\chi(x, \lambda_n) = k_n \phi(x, \lambda_n) \quad (6)$$

Consequently  $\phi(x, \lambda_n)$  and  $k_n \phi(x, \lambda_n)$  satisfy all of the boundary conditions (4).

It is also easy to show that  $\phi(x, \lambda_n)$  is an orthogonal set. For, in view of (2), (4) and (6) we have:

$$(\lambda_m - \lambda_n) \int_a^b \phi_m \phi_n dx = \int_a^b (\phi_m L \phi_n - \phi_n L \phi_m) dx = W(\phi_m, \phi_n) = 0.$$

$$(\lambda_m \neq \lambda_n)$$

Hence we can write:

$$\int_a^b \phi(x, \lambda_n) \phi(x, \lambda_m) dx = N^2 \delta_{mn} \quad (7)$$

where  $\delta_{mn}$  is the Kronecker delta (equal to zero for  $m \neq n$ , one for  $m=n$ )

It can be shown also that all roots of  $\omega(\lambda) = 0$  are real and distinct (see [4], pp 11-12).

From (5) and (6) it follows that:

$$k_n = \frac{\phi_0(b, \lambda_n) \cos \beta + \phi_{0,x}(b, \lambda_n) \sin \beta}{\phi_0(a, \lambda_n) \cos \alpha + \phi_{0,x}(a, \lambda_n) \sin \alpha}$$

$$= \frac{\chi_0(b, \lambda_n) \cos \beta + \chi_{0,x}(b, \lambda_n) \sin \beta}{\chi_0(a, \lambda_n) \cos \alpha + \chi_{0,x}(a, \lambda_n) \sin \alpha} \quad (8)$$

If now  $f(x)$  is an integrable function over  $(a, b)$  and if  $a < x < b$ , we have the Sturm-Liouville expansion of  $f(x)$ :

$$f(x) = \sum_{n=0}^{\infty} [k_n / \omega_{\lambda}(\lambda_n)] \phi(x, \lambda_n) \int_a^b \phi(y, \lambda_n) f(y) dy \quad (9)$$

For the convergence of (9) to  $f(x)$  see [4] pp 12-15 or [5] pp 275-276.

### 3. Finite Sturm-Liouville Transform

Let  $f(x)$  be a real continuous and integrable function of  $x$ , in the interval  $(a, b)$ . We define the finite Sturm-Liouville transform  $S\{f\} = F(\lambda_n)$  associated with system (2) and (4) by:

$$F(\lambda_n) = S\{f\} = \int_a^b f(y) \psi(y, \lambda_n) dy \quad (10)$$



where:

$$\psi(y, \lambda_n) = [k_n / \omega_{\lambda}(\lambda_n)]^{1/2} \phi(y, \lambda_n) \quad (11)$$

The inversion theorem now follows from (9):

$$f(x) = \sum_{n=0}^{\infty} F(\lambda_n) \psi(x, \lambda_n) \quad (12)$$

Here functions  $\psi(y, \lambda_n)$  are orthonormal, i.e.:

$$\int_a^b \psi(x, \lambda_n) \psi(x, \lambda_m) dx = \delta_{mn} \quad (13)$$

This is readily seen from (7) and (11).

Equations (10) and (12) are basic for the solution of some partial differential equations.

#### 4. Transform of Lf

We are now going to prove that:

$$\begin{aligned} S_x \{ Lf \} &= S_x \{ q(x)f - \partial^2 f / \partial x^2 \} = B_f(\lambda_n, t) \\ &+ \lambda_n F(\lambda_n, t) \end{aligned} \quad (14)$$

where:

$$\begin{aligned} B_f(\lambda_n, t) &= [k_n / \omega_{\lambda}(\lambda_n)]^{1/2} \left\{ -f(b, t) \frac{\cos \beta}{k_n} \right. \\ &\left. - f_{,x}(b, t) \frac{\sin \beta}{k_n} + f(a, t) \cos \alpha + f_{,x}(a, t) \sin \alpha \right\} \end{aligned} \quad (15)$$

is a function depending on the boundary values of  $f(x, t)$ ,  $f_{,x}(x, t)$  at  $x = a, b$ . Subscript  $f$  implies that the function  $B$  contains boundary values of the function  $f$ . Here  $S_x$  represents the finite Sturm-Liouville transform taken with respect to  $x$ :

$$S_x \{ \phi(x,t) \} = \bar{\phi}(\lambda_n, t) \quad (16)$$

To prove (14) we multiply  $Lf$  by  $\psi(x, \lambda_n)$  and integrate between  $(a, b)$ , that is, by definition we have:

$$S_x \{ Lf \} = \int_a^b [q(x)f(x,t) - \partial^2 f / \partial x^2] \psi(x, \lambda_n) dx \quad (17)$$

Integrating the second term in the integrand of (17) twice by part we obtain:

$$S_x \{ Lf \} = -[f_{,x} \psi - \psi_{,x} f]_a^b + \int_a^b f(x,t) L\psi(x, \lambda_n) dx \quad (18)$$

If we now use (4) and (6) in the first term of the right hand side and (2) inside the integrand we obtain (14).

##### 5. Solution of Some Partial Differential Equations

a) Let us consider a parabolic partial differential equation

$$Mv + p_4 v_{,t} = 0 \quad (19)$$

where  $Mv$  is defined by (1),  $p_i = p_i(y, t)$  are known and  $v(y, t)$  is the unknown dependent variable.

The use of transformation (3) reduces (19) to:

$$u_{,xx} + [\lambda - p(x, t)]u + (p_4/p_3)u_{,t} = 0 \quad (20)$$

where:

$$p(x, t) = q(x, t) - p_4 p_0 (k/2p_0)^2 \frac{\partial}{\partial t} (p_0/p_3 k^2) \quad (21)$$

Here  $q(x, t)$ ,  $k(t)$  and  $\lambda(t)$  are given by (3) with  $t$  as a parameter. If  $\phi(x, t, \lambda_n)$  and  $\psi(x, t, \lambda_n)$  are now obtained as before except with  $q(x)$  replaced by  $p(x, t)$ ,  $t$  a parameter,

and  $p_4/p_3$  a function of  $t$  alone, then (20), with the use of the finite Sturm-Liouville transform, can be reduced to an ordinary differential equation in  $t$ . For, let  $p_4/p_3 = r(t)$ , then application of the transform to (20) gives:

$$-B_u(\lambda_n, t) + (\lambda - \lambda_n)\bar{u}(\lambda_n, t) + r(t)\bar{u}_{,t}(\lambda_n, t) = 0 \quad (22)$$

Equation (22) is of first order. The complete solution is:

$$\begin{aligned} \bar{u}(\lambda_n, t) = & C(\lambda_n) \exp\left(\int^t \frac{\lambda_n - \lambda}{r} dt\right) + \left[\exp\left(\int^t \frac{\lambda_n - \lambda}{r} dt\right)\right] \cdot \\ & \int^t (B_u/r) \left[\exp\left(-\int^t \frac{\lambda_n - \lambda}{r} dt\right)\right] dt \end{aligned} \quad (23)$$

Now, inversion theorem (12) gives:

$$u(x, t) = \sum_{n=0}^{\infty} \bar{u}(\lambda_n, t) \psi_n(x, t, \lambda_n) \quad (24)$$

Therefore, the solution of (19) is effected. We must, however, remember that at least one of each pair of boundary values  $u(a, t)$ ,  $u_{,x}(a, t)$  and  $u(b, t)$ ,  $u_{,x}(b, t)$  must be known. By selecting  $\alpha$  and  $\beta$  properly in  $B_u(\lambda_n, t)$  we can make the terms containing the unknown pair zero. Of course any knowledge of the boundary values of  $u$  leading to the complete evaluation of  $B_u$  is sufficient for the solution.

b) A second order linear partial differential equation containing a parameter  $\lambda$  has the general form:

$$\begin{aligned} Nv + a_4 v_{,z} + a_5 v_{,\tau} + (a_6 + \lambda a_7) v &= 0 \\ Nv = a_1 v_{,zz} + 2a_2 v_{,z\tau} + a_3 v_{,\tau\tau} \end{aligned} \quad (25)$$

where  $a_1 = a_1(z, \tau)$ . If the equation is of elliptic type we have:

$$a_1 a_3 - a_2^2 > 0 \quad (26)$$

Consider now an elliptic equation (25). Let  $\xi(z, \tau) = \text{const.}$  and  $\eta(z, \tau) = \text{const.}$ , respectively be the solutions of the following differential equations:

$$\begin{aligned} d\tau/dz &= \zeta_1(z, \tau) & d\tau/dz &= \zeta_2(z, \tau) \\ \left. \begin{matrix} \zeta_1 \\ \zeta_2 \end{matrix} \right\} &= -\frac{a_2}{a_1} \pm 1 \left[ \frac{a_3}{a_1} - \left( \frac{a_2}{a_1} \right)^2 \right]^{1/2} \end{aligned} \quad (27)$$

If we set:

$$2y = \xi + \eta, \quad 2it = \xi - \eta \quad (28)$$

equation (25) can be transformed into the canonical form [6]:

$$v_{,yy} + p_1 v_{,y} + (p_2 + \lambda p_3) v + p_4 v_{,t} + v_{,tt} = 0 \quad (29)$$

where:

$$\begin{aligned} p_1 &= \frac{2}{\beta} p_y, \quad p_2 = 2a_6/\beta, \quad p_3 = 2a_7/\beta \\ p_4 &= \frac{2}{\beta} p_t, \quad \beta = a_1(y_{,z}^2 + t_{,z}^2) + 2a_2(y_{,z}y_{,\tau} + t_{,z}t_{,\tau}) \\ &\quad + a_3(y_{,\tau}^2 + t_{,\tau}^2) \end{aligned} \quad (30)$$

Here operator  $P$  is defined by:

$$P\phi = N\phi + a_4\phi_{,z} + a_5\phi_{,\tau} \quad (31)$$

and  $N\phi$  is given by the second of (25).

Using the transformation (3) with  $p_0 = 1$ , (29) becomes:

$$u_{,xx} + (\lambda - p)u + r_1 u_{,t} + r_0 u_{,tt} = 0 \quad (32)$$

where:

$$\begin{aligned} p &= q(x,t) - mp_4g_{,t} - mg_{,tt}, \quad r_0 = p_3^{-1} \\ r_1 &= mp_4g + 2mg_{,t}, \quad g = (p_3k^2)^{-1/4}, \\ m &= (k^2/p_3^3)^{1/4} \end{aligned} \quad (33)$$

In the special case where  $r_1$ , and  $r_0$  are functions of  $t$  alone equation (32) can be transformed into an ordinary differential equation by applying the finite Sturm-Liouville transform.

Hence:

$$r_0 \bar{u}_{,tt} + r_1 \bar{u}_{,t} + (\lambda - \lambda_n) \bar{u}(\lambda_n, t) = B_u(\lambda_n, t) \quad (34)$$

We can now either use transformation (3) with respect to  $t$ , with  $r_0$ ,  $r_1$  replacing  $p_0$ ,  $p_1$  and  $p_2 = 0$ ,  $p_3 = 1$  and find the solution of (34) and then invert it or we can solve (34) directly for  $\bar{u}(\lambda_n, t)$  and then (24) gives  $u(x, t)$ .

It must be remembered that this method has the advantage that it contains the boundary values of the function within the solution, consequently, the difficulty of satisfying boundary conditions is eliminated. The method is also valid in the cases where the usual technique of separation of variables fail.

## 6. Solutions $\phi_0$ and $\chi_0$ of Some Special Equations

Below are given two independent solutions  $\phi_0$  and  $\chi_0$  of some special differential equations (D. E.) of mathematical physics. Their canonical forms (2) will be given by only determining  $u$ ,  $q$ , and  $\lambda$ .

a) Bessel Functions

$$v_{,yy} + \frac{1}{y} v_{,y} + (s^2 - \frac{v^2}{y^2})v = 0 \quad (\text{D. E.})$$

$$u = x^{1/2}v, \quad y = x, \quad q = (v^2 - \frac{1}{4})x^{-2}, \quad \lambda = s^2$$

$$\phi_0(x, \lambda) = x^{1/2} J_v(xs), \quad \chi_0(x, \lambda) = x^{1/2} Y_v(xs)$$

Interval (a,b) should not contain  $x = 0$ .

b) Spherical Harmonics

$$[(1-y^2)v_{,y}]_{,y} + [v(v+1) - \frac{\mu^2}{1-y^2}]v = 0 \quad (\text{D. E.})$$

$$u = v \cos^{1/2}x, \quad y = \sin x, \quad q = \frac{\mu^2}{\cos^2 x} - \frac{1}{4} \tan^2 x - \frac{1}{2},$$

$$\lambda = v(v+1), \quad \phi_0(x, \lambda) = \cos^{1/2}x \cdot P_v^\mu(\sin x),$$

$$\chi_0(x, \lambda) = \cos^{1/2}x \cdot Q_v^\mu(\sin x)$$

Interval (a,b) should not contain  $(-1, 1)$ .

c) Hermite Polynomials

$$v_{,yy} - yv_{,y} + nv = 0 \quad (\text{D. E.})$$

$$u = e^{-x^2/4}v, \quad y = x, \quad q = -\frac{1}{2} + \frac{x^2}{4}, \quad \lambda = n,$$

$$\phi_0(x, \lambda) = e^{-x^2/4} \text{He}_n(x), \quad \chi_0(x, \lambda) = e^{-x^2/4} \text{he}_n(x)$$

d) Tschebyscheff Polynomials

$$(1-y^2)v_{,yy} - yv_{,y} + n^2v = 0 \quad (\text{D. E.})$$

$$u = v, \quad y = \sin x, \quad q = 0, \quad \lambda = n^2$$

$$\phi_0(x, \lambda) = T_n(\sin x) = \cos n(\sin^{-1}y),$$

$$\chi_0(x, \lambda) = U_n(\sin x) = \sin n(\sin^{-1}y)$$

e) Mathieu Functions

$$u_{,xx} + (\lambda - 2h^2 \cos 2x)u = 0 \quad (\text{D. E.})$$

$$\phi_0(x, \lambda) = ce_{2n}(x), \quad ce_{2n+1}(x), \quad se_{2n}(x), \quad s_{2n+1}(x)$$

$$\chi_0(x, \lambda) = ce_{2n}^{(2)}(x), \quad ce_{2n+1}^{(2)}(x), \quad se_{2n}^{(2)}(x)$$

$$s_{2n+1}^{(2)}(x)$$

( $n = 0, 1, 2, \dots$  except for  $se_{2n}$  only  $n = 1, 2, \dots$ )

$\phi_0(x, \lambda)$  is periodic.

f) Wittaker Functions

$$v_{,yy} + \left(-\frac{1}{4} + \frac{k}{y} + \frac{\frac{1}{4} - \mu^2}{y^2}\right)v = 0 \quad (\text{D. E.})$$

$$u = (x/2)^{1/2}v, \quad y = x^2/4, \quad q = \frac{16\mu^2 - 17}{4x^2} + \frac{x^2}{16}$$

$$\phi_0(x, \lambda) = (x/2)^{1/2} W_{k, \mu}(x^2/4), \quad \chi_0(x, \lambda) = (x/2)^{1/2}$$

$$W_{-k, \mu}(-x^2/4) \quad (*)$$

Interval (a, b) should not contain  $x = 0$ .

7. Application

As an illustration we solve a heat conduction problem which may be looked upon as a mathematical model of volcanos. The problem also has application in the exhaust parts of jet engines. As far as I know the problem has not previously been solved.

---

\* Here comma does not mean differentiation.

Consider the conical shell enclosed by two coaxial cones having the same apex  $O$ , and two concentric spheres having  $O$  as their center (Fig. 1).




Fig. 1

In polar coordinates the inner and outer conical surfaces are given by  $\theta = \theta_0$  and  $\theta = \theta_1$ , and the end surfaces by  $r = r_0$  and  $r = r_1$ .

Problem: Determine the steady temperature distribution within the shell under the general axi-symmetric boundary conditions in temperature. This is the general Dirichlet problem for the domain under consideration. The differential equation and boundary conditions (B. C.) are given below:



$$\begin{aligned} \text{(D. E.) } \Delta V &= r^{-2}(r^2 V_{,r})_{,r} + r^{-2}[(1-y^2)V_{,y}]_{,y} = 0 \\ y &= \cos \theta, \quad r_0 < r < r_1, \quad \theta_0 < \theta < \theta_1 \end{aligned} \quad (35)$$

$$\begin{aligned} \text{(B. C.) } V &= V_0(r) \quad \text{for } \theta = \theta_0 \\ V &= V_1(r) \quad \text{for } \theta = \theta_1 \end{aligned} \left. \vphantom{\begin{aligned} V &= V_0(r) \\ V &= V_1(r) \end{aligned}} \right\} r_0 < r < r_1 \\ \\ V &= V_3(\theta) \quad \text{for } r = r_0 \\ V &= V_4(\theta) \quad \text{for } r = r_1 \end{aligned} \left. \vphantom{\begin{aligned} V &= V_3(\theta) \\ V &= V_4(\theta) \end{aligned}} \right\} \theta_0 < \theta < \theta_1 \quad (36)$$

Here  $V(r, y)$  is the temperature function,  $r$  and  $\theta$  are the polar coordinates.

Solution: We can exclude the second term of (D. E.) (35) if we use a finite Sturm-Liouville transform associated with the Legendre equation:

$$[(1-y^2)v_{,y}]_{,y} + v(v+1)v = 0 \quad (37)$$

In view of (6, b) with  $\mu = 0$  we find that if we select  $y = \sin x$ ,  $u = v \cos^{1/2} x$  we can transform (37) to canonical form leading to solutions:

$$\begin{aligned} \phi_0(x, \lambda) &= \cos^{1/2} x \cdot P_v(\sin x), \quad \chi_0(x, \lambda) = \cos^{1/2} x \\ &\cdot Q_v(\sin x), \quad \lambda = v(v+1) \end{aligned} \quad (38)$$

where  $P_v$  and  $Q_v$  are Legendre functions of the first and second kind, respectively. In view of (14), when we apply the transform to (35) the second term gives  $-v_n(v_n+1) - B_v(r, \lambda_n)$ . Now  $B_v$  contains four arbitrary functions which must be specified on the surfaces  $\theta = \theta_0$  and  $\theta = \theta_1$ . The terms containing the derivatives of  $v$  are not given. Thus if we

select  $\alpha = \beta = 0$  these terms drop out, leaving the terms containing  $v_0(r)$  and  $v_1(r)$ , which are given. Hence  $\phi(x, \lambda)$  and  $\omega_0(\lambda)$  of (5) become:

$$\begin{aligned} \phi(x, \lambda) &= \cos^{1/2} x_0 \cos^{1/2} x [P_v(\sin x) Q_v(\sin x_0) \\ &\quad - P_v(\sin x_0) Q_v(\sin x)] , \quad x_0 = \frac{\pi}{2} - \theta_0 , \\ x_1 &= \frac{\pi}{2} - \theta_1 , \quad \omega_0(\lambda) = 1 \end{aligned} \quad (39)$$

By (11) and (8) we have:

$$\begin{aligned} \psi(x, \lambda_n) &= [k_n / \omega_{\lambda}(\lambda_n)]^{1/2} \phi(x, \lambda_n) , \\ k_n &= \frac{\cos^{1/2} x_1 P_{v_n}(\sin x_1)}{\cos^{1/2} x_0 P_{v_n}(\sin x_0)} \end{aligned} \quad (40)$$

Calculation of  $\omega(\lambda) = \phi \chi_x - \chi \phi_x$  gives

$$\begin{aligned} \omega(\lambda) &= \cos^{1/2} x_0 \cos^{1/2} x_1 [P_v(\sin x_0) Q_v(\sin x_1) \\ &\quad - P_v(\sin x_1) Q_v(\sin x_0)] \end{aligned} \quad (41)$$

Consequently, the roots  $\lambda_n = v_n(v_n+1)$  of  $\omega(\lambda) = 0$  satisfy the following equation

$$\frac{P_v(\sin x_1)}{P_v(\sin x_0)} = \frac{Q_v(\sin x_1)}{Q_v(\sin x_0)} \quad (42)$$

Therefore  $\psi(x, \lambda_n)$  is completely determined.

After transforming (35) with  $y = \sin x$  and  $u = v \cos^{1/2} x$  we apply the finite Sturm-Liouville transform. That is, we multiply the equation by  $\psi(x, \lambda_n)$  and integrate between  $x_0$  and  $x_1$ . The result is:

$$(r^2 \bar{V}_{,r})_{,r} - v_n(v_n+1) \bar{V} = B_V(r, \lambda_n) \quad (43)$$

$$B_V(r, \lambda_n) = [k_n / \omega_{\lambda}(\lambda_n)]^{1/2} [V_0(r) - k_n^{-1} V_1(r)]$$

Equation (43) is an ordinary differential equation of Euler type whose solution can be found by variation of parameters.

Hence:

$$\begin{aligned} \bar{V}(r, v_n) &= C_1(v_n) r^{v_n} + C_2(v_n) r^{-v_n-1} + F(r, v_n) \\ F(r, v_n) &= \int^r \frac{B_V(\rho, \lambda_n)}{1+2v_n} (r^{v_n} \rho^{-v_n-1} - r^{-v_n-1} \rho^{v_n}) d\rho \end{aligned} \quad (44)$$

Let the transforms of  $V_3(\theta)$  and  $V_4(\theta)$  be  $\bar{V}_3(v_n)$  and  $\bar{V}_4(v_n)$ .

$C_1$  and  $C_2$  will then be determined from the remaining conditions:

$$\begin{aligned} \bar{V} &= \bar{V}_3(v_n) \quad , \quad \text{for } r = r_0 \\ \bar{V} &= \bar{V}_4(v_n) \quad , \quad \text{for } r = r_1 \end{aligned} \quad \theta_0 < \theta < \theta_1$$

This gives two linear equations for  $C_1$  and  $C_2$  whose solutions are:

$$\begin{aligned} C_1(v_n) &= \frac{[\bar{V}_3(v_n) - F(r_0, v_n)] r_1^{-v_n-1} - [\bar{V}_4(v_n) - F(r_1, v_n)] r_0^{-v_n-1}}{r_0^{v_n} r_1^{-v_n-1} - r_1^{v_n} r_0^{-v_n-1}} \\ C_2(v_n) &= \frac{[\bar{V}_4(v_n) - F(r_1, v_n)] r_0^{v_n} - [\bar{V}_3(v_n) - F(r_0, v_n)] r_1^{v_n}}{r_0^{v_n} r_1^{-v_n-1} - r_1^{v_n} r_0^{-v_n-1}} \end{aligned} \quad (45)$$

Hence,  $\bar{V}(r, v_n)$  is completely determined. The inversion theorem (12) now gives:

$$V(r, x) = \sum_n \bar{V}(r, v_n) \psi(x, \lambda_n) \quad (46)$$

where the summation is extended over all roots of (42).

The analysis given above is formal. It can, however, be made rigorous by showing that the solution satisfies both the differential equation and the boundary conditions.

It may be worth while to remark that with the use of the method of separation of variables the solution of the above problem would have been difficult and lengthy, if not impossible.

References

1. Churchill, R. V., Modern Operational Mathematics in Engineering, McGraw-Hill Book Company, 1944, p. 267-291.
2. Tranter, C. J., "Legendre Transform", Quarterly Journal of Mathematics, (Oxford), (2), 1, 1950, 1-8.
3. Scott, E. J., "Jacobi Transform", Quarterly Journal of Mathematics, (Oxford), (4), 13, 1953, 36-40.
4. Titchmarsh, E. C., Eigenfunction Expansion Associated with Second Order Differential Equations, Oxford University Press, 1946.
5. Ince, E. L., Ordinary Differential Equations, Dover Publications, 1944.
6. Tamarkin, J. D. and Feller, W., Partial Differential Equations, Brown University Notes, June 23-Sept. 13, 1941. 1-5.

Contributing Personnel:

Dr. A. C. Eringen, Research Associate Professor  
of Mechanics

Mr. T. King, Staff Computer

Respectfully submitted,

*A. C. Eringen*  
\_\_\_\_\_  
A. C. Eringen

*William R. Osgood*  
\_\_\_\_\_  
William R. Osgood  
Chairman, Department  
of Mechanics

## Distribution List

### I: Administrative, Reference and Liaison Activities of ONR

Chief of Naval Research Department of the Navy Washington 25, D. C. Attn: Code 438 (2) Code 432 (1) Code 466(via Code 108) (1)	Library of Congress Washington 25, D. C. Attn: Navy Research Section (2)
Director, Naval Research Lab. Washington 25, D. C. Attn: Tech. Info. Officer (9) Technical Library (1) Mechanics Division (2)	Office of Naval Research Branch Office The John Crerar Library Building Tenth Floor, 86 E. Randolph St. Chicago 1, Illinois (2)

### II: Department of Defense and other interested Government Activities

#### (A) GENERAL

Research and Development Board  
Department of Defense  
Pentagon Building  
Washington 25, D. C.  
Attn: Library(Code 3D-1075) (1)

Armed Forces Special Weapons  
Project  
P.O. Box 2610  
Washington, D. C.  
Attn: Col. G.F.Blunda (2)

Joint Task Force 3  
12 St. and Const. Ave.,N.W.  
(Temp. U)  
Washington 25, D. C.  
Attn: Major B. D. Jones (1)

#### (B) ARMY

Chief of Staff  
Department of the Army  
Research and Development Div.  
Washington 25, D. C.  
Attn: Chief of Res. and Dev. (1)

Office of the Chief of Engineers  
Assistant Chief for Works  
Department of the Army  
Bldg. T-7, Gravelly Point  
Washington 25, D. C.  
Attn: Structural Branch  
(R. L. Bloor) (1)

Engineering Research and  
Development Laboratory  
Fort Belvoir, Virginia  
Attn: Structures Branch (1)

# Distribution List (Cont.)

Office of the Chief of Engineers  
Asst. Chief for Military  
Construction  
Department of the Army  
Bldg. T-3, Gravelly Point  
Washington 25, D. C.  
Attn: Structures Branch  
(M. F. Carey) (1)  
Protective Construction  
Branch (I.O.Thorley) (1)

Office of the Chief of Engineers  
Asst. Chief for Military  
Operations  
Department of the Army  
Bldg T-7, Gravelly Point  
Washington 25, D. C.  
Attn: Structures Development  
Branch (W.F.Woollard) (1)

U. S. Army Waterways Experiment  
Station  
P. O. Box 631  
Halls Ferry Road  
Vicksburg, Mississippi  
Attn: Col. H. J. Skidmore (1)

The Commanding General  
Sandia Base, P.O.Box 5100  
Albuquerque, New Mexico  
Attn: Col. Canterbury (1)

Operations Research Officer  
Department of the Army  
Ft. Lesley J. McNair  
Washington 25, D. C.  
Attn: Howard Brackney (1)

Office of Chief of Ordnance  
Office of Ordnance Research  
Department of the Army  
The Pentagon Annex No. 2  
Washington 25, D. C.  
Attn: ORDTB-PS (1)

Ballistics Research Laboratory  
Aberdeen Proving Ground  
Aberdeen, Maryland  
Attn: Dr. C. W. Lampson (1)

## (C) NAVY

Chief of Naval Operations  
Department of the Navy  
Washington 25, D. C.  
Attn: OP-31 (1)  
OP-363 (1)

Chief of Bureau of Ships  
Department of the Navy  
Washington 25, D. C.  
Attn: Director of Research (2)  
Code 423 (1)  
Code 442 (1)  
Code 421 (1)

Director, David Taylor Model Basin  
Department of the Navy  
Washington 7, D. C.  
Attn: Code 720, Structures  
Division (1)  
Code 740, Hi-Speed  
Dynamics Div. (1)

Commanding Officer  
Underwater Explosion Research Div.  
Code 290  
Norfolk Naval Shipyard  
Portsmouth, Virginia (1)

Commander  
Portsmouth Naval Shipyard  
Portsmouth, N. H.  
Attn: Design Division (1)

Director, Materials Laboratory  
New York Naval Shipyard  
Brooklyn 1, New York (1)

Chief of Bureau of Ordnance  
Department of the Navy  
Washington 25, D. C.  
Attn: Ad-3, Technical Library (1)  
Roc, P.H. Girouard (1)

Naval Ordnance Laboratory  
White Oak, Maryland  
RFD 1, Silver Spring, Maryland  
Attn: Mechanics Division (1)  
Explosive Division (1)  
Mech. Evaluation Div. (1)



Distribution List (Cont.)

(C) NAVY

Commander  
U.S. Naval Ordnance Test Station  
Inyokern, California  
Post Office - China Lake, Calif.  
Attn: Scientific Officer (1)

Naval Ordnance Test Station  
Underwater Ordnance Division  
Pasadena, California  
Attn: Structures Division (1)

Chief of the Bureau of Aeronautics  
Department of the Navy  
Washington 25, D. C.  
Attn: TD-41, Technical Library (1)

Chief of Bureau of Ships  
Department of the Navy  
Washington 25, D. C.  
Attn: Code P-314 (1)  
Code C-313 (1)

Officer in Charge  
Naval Civil Engr. Research and  
Evaluation Laboratory  
Naval Station  
Port Hueneme, California (1)

Superintendent  
U.S. Naval Postgraduate School  
Annapolis, Maryland (1)

(D) AIR FORCES

Commanding General  
U.S. Air Forces  
The Pentagon  
Washington 25, D. C.  
Attn: Res. and Develop. Div. (1)

Deputy Chief of Staff, Operations  
Air Targets Division  
Headquarters, U.S. Air Forces  
Washington 25, D. C.  
Attn: AFOIN-T/PV (1)

Flight Research Laboratory  
Wright-Patterson Air Force Base  
Dayton, Ohio  
Attn: Chief, Applied Mechanics  
Group (1)

(E) OTHER GOVERNMENT AGENCIES

U.S. Atomic Energy Commission  
Division of Research  
Washington, D. C. (1)

Director, National Bureau of  
Standards  
Washington, D. C.  
Attn: Dr. W. H. Ramberg (1)

Forest Products Laboratory  
Madison 5, Wisconsin (1)

Supplementary Distribution List

Professor Lynn Beedle  
Fritz Engineering Laboratory  
Lehigh University  
Bethlehem, Pennsylvania (1)

Professor R. L. Bisplinghoff  
Dept. of Aeronautical Engineering  
Massachusetts Institute of Tech.  
Cambridge 39, Massachusetts (1)

Professor G. F. Carrier  
Graduate Division of Applied  
Mathematics  
Brown University  
Providence, R. I. (1)

Professor R. J. Dolan  
Dept. of Theoretical and Applied  
Mechanics  
University of Illinois  
Urbana, Illinois (1)

# Distribution List (Cont.)

Professor Hans Bleich Dept. of Civil Engineering Columbia University Broadway at 117th St. New York 27, New York	(1)	Professor L. S. Jacobsen Department of Mechanical Eng. Stanford University Stanford, California	(1)
Professor B. A. Boley Dept. of Aeronautical Eng. Ohio State University Columbus, Ohio	(1)	Dr. W. H. Hoppmann Department of Applied Mechanics Johns Hopkins University Baltimore, Maryland	(1)
Professor Lloyd Donnell Department of Mechanics Illinois Institute of Tech. Technology Center Chicago 16, Illinois	(1)	Professor George Lee Department of Mathematics Rensselaer Polytechnic Institute Troy, New York	(1)
Professor B. Fried Dept. of Mechanical Engineering Washington State College Pullman, Washington	(1)	Professor Glen Murphy, Head Department of Theoretical and Applied Mechanics Iowa State College Ames, Iowa	(1)
Mr. Martin Goland Midwest Research Institute 4049 Pennsylvania Avenue Kansas City 2, Missouri	(1)	Professor N. M. Newmark Department of Civil Engineering University of Illinois Urbana, Illinois	(1)
Dr. J. N. Goodier School of Engineering Stanford University Stanford, California	(1)	Professor Jesse Ormondroyd University of Michigan Ann Arbor, Michigan	(1)
Professor M. Hetenyi Walter P. Murphy Professor Northwestern University Evanston, Illinois	(1)	Dr. R. P. Peterson, Director Applied Physics Division Sandia Laboratory Albuquerque, New Mexico	(1)
Dr. N. J. Hoff, Head Department of Aeronautical Engineer- ing and Applied Mechanics Polytechnic Institute of Brooklyn 99 Livingston Street Brooklyn 2, New York	(1)	Dr. A. Phillips School of Engineering Stanford University Stanford, California	(1)
Dr. J. H. Hollomon General Electric Research Laboratories 1 River Road Schenectady, New York	(1)	Dr. W. Prager, Chairman Graduate Division of Applied Mathematics Brown University Providence 12, R. I.	(1)

Distribution List (Cont.)

Dr. S. Raynor Armour Research Foundation Illinois Institute of Technology Chicago 16, Illinois	(1)	Professor E. Sternberg Illinois Institute of Technology Technology Center Chicago 16, Illinois	(1)
Professor E. Reissner Department of Mathematics Massachusetts Institute of Tech. Cambridge 39, Massachusetts	(1)	Professor F. K. Teichmann Department of Aeronautical Engineering New York University University Heights, Bronx New York, N. Y.	(1)
Professor M. A. Sadowsky Illinois Institute of Technology Technology Center Chicago 16, Illinois	(1)	Professor C. T. Wang Department of Aeronautical Engineering New York University University Heights, Bronx New York, N. Y.	(1)
Professor Paul Lieber Department of Aeronautical Eng. Rensselaer Polytechnic Institute Troy, New York	(1)	Project File	(2)
Professor J. E. Stallmeyer Talbot Laboratory Department of Civil Engineering University of Illinois Urbana, Illinois	(1)	Project Staff	(5)
		For possible future distribution by the University	(10)
Dr. C. B. Smith College of Arts and Sciences Department of Mathematics Walker Hall University of Florida Gainesville, Florida	(1)		
Professor J. R. Andersen Towne School of Engineering University of Pennsylvania Philadelphia, Pennsylvania	(1)		
Commander U.S. Naval Proving Ground Dahlgren, Virginia	(1)		